

CALCULATION OF TOPOLOGICAL CHARGE OF REAL FINITE-GAP SINE-GORDON SOLUTIONS USING THETA-FUNCTIONAL FORMULAE

P. G. GRINEVICH AND K. V. KAIPA

The most basic characteristic of real solutions $u(x, t)$ of the sine-Gordon equation $u_{tt} - u_{xx} + \sin u = 0$ which are quasiperiodic in the x -variable is the *density of topological charge* defined as:

$$\bar{n} = \lim_{T \rightarrow \infty} \frac{u(x+T, t) - u(x, t)}{2\pi T}$$

The real finite-gap solutions $u(x, t)$ are expressed in terms of the Riemann theta function of non-singular hyperelliptic curves $\Gamma : \mu^2 = \lambda \prod_{i=1}^{2g} (\lambda - E_i)$ and a generic positive divisor D of degree g on Γ , where the spectral data (Γ, D) must satisfy some reality conditions. The problem of calculating $\bar{n}$ in terms of the spectral data was first studied in [3] and [6]. A solution to this problem was obtained in [4] avoiding the use of the θ -functional formula for $u(x, t)$. As pointed out by S.P. Novikov, if the θ -functional form of solutions is to be considered as an effective one, then it should be possible to calculate $\bar{n}$ directly from the θ -functional formulae. We achieve this goal in the present note, using a new *multiscale* or *elliptic limit* of real finite-gap sine-Gordon solutions. The authors would like to express their gratitude to Professor Novikov for attracting their attention to this problem and stimulating discussions.

The reality condition on Γ is that $\{E_1, \dots, E_{2g}\} = \{\overline{E_1}, \dots, \overline{E_{2g}}\}$ for all i , and $E_i < 0$ if $E_i \in \mathbb{R}$. The reality condition on the divisor found by Cherednik in [1] is: $D + \tau D - 0 - \infty = \mathcal{K}$ where $\tau(\lambda, \mu) = (\bar{\lambda}, \bar{\mu})$ and $\mathcal{K}$ is the canonical class. Such a divisor will be called *admissible*. In order to write the θ -functional formula for $u(x, t)$ we need some notation. Let a_i, b_j for $1 \leq i, j \leq g$ be a symplectic basis of cycles on Γ and $\vec{\omega} = (\omega_1, \dots, \omega_g)$ holomorphic differentials satisfying $\int_{a_j} \omega_i = \delta_{ij}$. The Riemann matrix of Γ is the matrix defined by $B_{ij} = \int_{b_j} \omega_i$. The Abel-Jacobi map $A : \Gamma \rightarrow J(\Gamma)$ is defined by $P \mapsto \int_{\infty}^P \vec{\omega}$, where $J(\Gamma) = \mathbb{C}^g / \{\mathbb{Z}^g + B\mathbb{Z}^g\}$ is the Jacobian variety of Γ . Let $K \in J(\Gamma)$ be the associated vector of Riemann constants. Let $A(0) \equiv \epsilon'/2 + B\epsilon/2$, for some $\epsilon, \epsilon' \in \mathbb{Z}^g$. We define $U = (\frac{\vec{\omega}}{d\sqrt{\lambda-1}})(\infty)$ and $V = (\frac{\vec{\omega}}{d\sqrt{\lambda}})(0)$. The θ -functional formula for $e^{iu(x,t)}$ can be written as follows (see [5], [2]):

$$(1) \quad e^{iu(x,t)} = C_1 \frac{\theta(A(0) + z(x, t)) \theta(-A(0) + z(x, t))}{\theta^2(z(x, t))}$$

where $z(x, t) = -A(D) + ix(V - U)/4 - it(U + V)/4 - K$, $C_1^2 = \exp(\pi i \epsilon^t B \epsilon / 2)$, and

$$\theta(z|B) = \sum_{n \in \mathbb{Z}^g} \exp(\pi i n^t B n) \exp(2\pi i n^t z) \quad \text{for } z \in \mathbb{C}^g$$

In order to compute $\bar{n}$ for real solutions, we choose a special basis of cycles a_i, b_j (figure (1) with parameter $k = 1$) as suggested in [4]. Here $E_1, \dots, E_{2m}$ are the non-zero real branch points and $E_{2j} = \overline{E_{2j-1}}$ for $m+1 \leq j \leq g$. This basis satisfies $\tau a_i = -a_i$ for all i , $\tau b_i = b_i$

for $1 \leq i \leq m$ and $\tau b_i = b_i + a_i$ for $m+1 \leq i \leq g$, therefore $\text{Re}(B) = -1/2 \begin{pmatrix} 0 & 0 \\ 0 & I_{g-m} \end{pmatrix}$ where I_{g-m} is the identity matrix of size $g-m$. (From here on, all vectors v written as $(v_1, v_2)^t$ are understood to be split into blocks of length m and $g-m$). We have $A(0) = \epsilon'/2 + B\epsilon/2$ with $\epsilon' = (0, 1)^t$ and $\epsilon = (1, 0)^t$. The reality condition on the divisor becomes

$$(2) \quad A(D) = x + B \begin{pmatrix} 1/2 - s/4 \\ 1/2 \end{pmatrix}, \quad s = \begin{pmatrix} s_1 \\ s_2 \\ \vdots \\ s_m \end{pmatrix}$$

for some $x \in \mathbb{R}^g / \mathbb{Z}^g$ and for some $s_k \in \{-1, 1\}$

Thus the set $\{z \in J(\Gamma) \mid z = -A(D) - K \text{ with } D \text{ admissible}\}$ has 2^m components characterized by the symbol $\vec{s} = (s_1, s_2, \dots, s_m)$, each of which is a real g -dimensional torus denoted by $T_{\vec{s}}$. The vectors iU, iV are purely real and hence $z(x, t) \in T_{\vec{s}}$. It is a consequence of the Cherednik reality condition that the $T_{\vec{s}}$ are disjoint from $\Theta \cup (\Theta - A(0))$ where $\Theta = \{z \in J(\Gamma) \mid \theta(z) = 0\}$. Therefore the real solutions $u(x, t)$ given by equation (1) are non-singular. We also define functions $u_j(T)$ for $T \in [0, 1]$ by the formula:

$$e^{iu_j(T)} = C_1 \frac{\theta(A(0) + z_j(T)) \theta(-A(0) + z_j(T))}{\theta^2(z_j(T))}, \quad z_j(T) = -A(D) - Te_j - K$$

where e_j is the j -th standard basis vector of $\mathbb{C}^g$, and therefore $z_j(T)$ represents the j -th basic cycle of $T_{\vec{s}}$. The topological charge density $\bar{n}$ is given by the formula (see [4]):

$$\bar{n} = \sum_{j=1}^g (iU_j - iV_j) n_j / 4, \quad \text{where} \quad n_j = \frac{1}{2\pi i} \int_{T=0}^{T=1} d \log e^{iu_j(T)}$$

It is convenient to replace $\epsilon = (1, 0)^t$ in the expression $A(0) = \epsilon'/2 + B\epsilon/2$ with $\tilde{\epsilon}$ defined by $\tilde{\epsilon}_j = (-1)^j s_j$ for $1 \leq j \leq m$ and $\tilde{\epsilon}_j = 0$ for $j > m$. Using the transformation rule $\theta(z + N + BM) = \theta(z) \exp(-2\pi i M^t z - \pi i M^t B M)$ we get

$$(3) \quad n_j = -\tilde{\epsilon}_j + \frac{2}{2\pi i} \int_{T=0}^1 d \log \left(\frac{\theta(B\tilde{\epsilon}/2 + z_j(T))}{\theta(z_j(T))} \right)$$

where we have also used the fact that the cycles $\epsilon'/2 + B\tilde{\epsilon}/2 + z_j(T)$ and $B\tilde{\epsilon}/2 + z_j(T)$ are homologous in the real torus $T_{-\vec{s}}$. The charges n_j are easily calculated using formula (3) in two special cases:

Lemma 1. *In the case when $m = 0$ (no real branch points) we have $n_j = 0$ for all j . In the case $g = m = 1$, we have $n_1 = s_1$. For later use, we also calculate n_1 from (3) if K is taken to be $1/2$ instead of the correct value $(1 + \tau)/2$, and s_1 is replaced by $-s_1$. In this case $n_1 = -s_1$.*

We reduce the calculation of the charges n_j in the general case to the two special cases of Lemma 1. We consider the following family of real nonsingular hyperelliptic curves $\Gamma(k)$ depending on the real parameter $k \in [1, \infty)$:

$$\Gamma(k) : \mu^2 = \lambda \prod_{i=1}^m (\lambda - k^{i-1} E_{2i-1}) (\lambda - k^{i-1} E_{2i}) \prod_{i=2m+1}^{2g} (\lambda - k^m E_i)$$

The basic cycles $a_i(k), b_j(k)$ on $\Gamma(k)$ are chosen as shown in figure (1). The original curve Γ coincides with $\Gamma(1)$. Let $\mathcal{C}_i$ for $1 \leq i \leq m$ be the real elliptic curves defined by $\mathcal{C}_i : y^2 = x(x - E_{2i-1})(x - E_{2i})$ and let $\mathcal{C}_{m+1}$ be the real hyperelliptic curve $\mathcal{C}_{m+1} : y^2 = x \prod_{i=2m+1}^{2g} (\lambda - E_i)$.

Let $B(k)$ denote the Riemann matrix of the curve $\Gamma(k)$ with respect to the basic cycles $a_i(k), b_j(k)$.




FIGURE 1. Basic cycles and cuts on $\Gamma(k)$ for $g = 4, m = 2$

Theorem 1. *The limiting curve $\Gamma(\infty) = \lim_{k \rightarrow \infty} \Gamma(k)$ is a nodal curve with m nodes when $m < g$, and $m - 1$ nodes when $m = g$. The irreducible components of the normalization of $\Gamma(\infty)$ are $\{\mathcal{C}_i \mid 1 \leq i \leq m + 1\}$ when $m < g$, and $\{\mathcal{C}_i \mid 1 \leq i \leq g\}$ when $m = g$. The limiting Riemann matrix $B(\infty) = \lim_{k \rightarrow \infty} B(k)$ is block diagonal $B(\infty) = \text{diag}(B_1, B_2)$ where $B_1 = \text{diag}(\tau_1, \dots, \tau_m)$ for some purely imaginary τ_j in the complex upper half plane, and $\text{Re}(B_2) = -1/2 I_{g-m}$.*

The charges n_j being integers are constant during the deformation $k : 1 \rightarrow \infty$. Therefore n_j can be calculated from formula (3) using the Riemann matrix $B(\infty)$.

Since $\theta((z_1, z_2)^t \mid \text{diag}(B_1, B_2)) = \theta(z_1 \mid B_1) \theta(z_2 \mid B_2)$, using Lemma 1 together with the formula $K = \sum_{i=1}^g A(E_{2i-1}) = \frac{1}{2} \binom{1}{\nu_2} + \frac{1}{2} B \binom{\nu_1}{1}$ where $(\nu_1, \nu_2)^t = (1, 2, \dots, g)^t$, we obtain the result:

$$(4) \quad n_j = \begin{cases} (-1)^{j-1} s_j & \text{if } 1 \leq j \leq m \\ 0 & \text{if } j > m \end{cases}$$

The details will appear in a future article.

Remarks

- (1) In [4], the admissible divisors were characterized by certain symbols $\{s'_1, \dots, s'_m\} \in \{\pm 1\}^m$ defined as follows. Given an admissible divisor $D = \{(\lambda_i, \mu_i) \mid 1 \leq i \leq g\}$ let $P(\lambda)$ be the unique polynomial of degree $g - 1$ interpolating the g points $(\lambda_i, \mu_i/\lambda_i)$. Then $P(\lambda)$ is real and s'_j is defined to be the sign of $P(\lambda)$ over $[E_{2j}, E_{2j-1}]$. It was shown in [4] that the charges n_j are equal to $(-1)^{j-1} s'_j$ for $j \leq m$ and $n_j = 0$ for $j > m$. Comparing with formula (4), it follows that the symbols s'_j and s_j coincide for all j .

- (2) The multiscale limit of the spectral curve constructed above was used only for a topological argument. The sine-Gordon solutions $u(x, t, k)$ associated with the spectral curve $\Gamma(k)$ (and admissible divisors $D(k)$) depend on the vectors $U(k)$ and $V(k)$ mentioned in the introduction. As $k \rightarrow \infty$, some component of $U(k)$ will diverge to ∞ . Thus there is no limiting solution. However asymptotic expansion in the parameter k of $u(x, t, k)$ involving elliptic (genus 1) solutions can be written. This will be investigated in a future work.

REFERENCES

- [1] I.V. Cherednik, *Reality conditions in “finite-zone” integration*, Sov. Phys. Dokl. **25** (1980), 450–452.
- [2] B.A. Dubrovin, S.M. Natanzon, *Real two-zone solutions of the sine-Gordon equation*, Funct. Anal. Appl. **16** (1982), 21–33.
- [3] B.A. Dubrovin, S.P. Novikov, *Algebro-geometrical Poisson brackets for real finite-zone solutions of the Sine-Gordon equation and the nonlinear Schrödinger equation*, Sov. Math. Dokl. **26** (1982), no. 3, 760–765.
- [4] P.G. Grinevich, S.P. Novikov, *Topological charge of the real periodic finite-gap sine-Gordon solutions*, Comm. Pure Appl. Math. **56** (2003), no. 7 956–978.
- [5] V.A. Kozel, V.P. Kotlyarov, *Almost periodic solutions of the equation $u_{tt} - u_{xx} + \sin u = 0$* . (Russian), Dokl. Akad. Nauk Ukrain. SSR Ser. A (1976), no. 10 878–881.
- [6] S.P. Novikov, *Algebrotopological approach to the reality problems. Real action variables in the theory of finite-zone solutions of the Sine-Gordon equation*, Zap. Nauchn. Sem. LOMI: Differential geometry, Lie groups and mechanics. VI, **133** (1984), 177–196.

L.D. LANDAU INSTITUTE FOR THEORETICAL PHYSICS, RUSSIAN ACADEMY OF SCIENCES
E-mail address: pgg@landau.ac.ru

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK
E-mail address: kaipa@math.umd.edu